

POSITION STATEMENT

POSITION STATEMENT ON HOME BIRTH

RATIONALE

The best available evidence from North America and international studies demonstrates that midwife attended home births are associated with optimal labour and birth outcomes. Midwife attended planned home births in Canada are associated with fewer obstetrical interventions and no increase in maternal/fetal/neonatal mortality or morbidity compared to births planned to be in hospital.

The Canadian Association of Midwives (CAM/ACSF) recognizes that planned home birth for healthy women is safe and that the definition of safety in maternity care includes physical safety, as well as a woman's self-defined values of cultural, spiritual and emotional safety for herself, her baby and her family.

STATEMENT OF BELIEF

- Women have the right to make an informed choice regarding their place of birth.
- Planned home birth should be recommended for healthy, low-risk women.
- Women who plan home births experience reduced obstetric and neonatal interventions and a course of care that facilitates normal, physiologic birth.
- Evidence-based birth site selection requires ongoing assessment and care throughout the
 antepartum, intrapartum and postpartum periods. Midwives have unique expertise in the use of
 low technology procedures and are skilled in intrapartum assessment and care in low resource
 settings. Therefore, midwives are ideal primary care providers for mothers and newborns in the
 home setting.
- Integration of midwives into a health care system that allows for timely consultation, referral, and transfer when necessary, enhances the safety of planned home birth.
- Women have the right to refuse recommended care. If a woman will not accept her midwife's
 recommendation for transfer to hospital from planned home birth after she begins labour, it is a
 midwife's ethical duty to remain with the woman and offer her midwifery skills in the home setting
 to the degree that the woman will accept care.

POSITION STATEMENT

To ensure women's choice and safe access to planned home birth, CAM/ACSF calls on every province and territory to expand regulated and funded midwifery services.

Midwives in Canada should provide all women with informed choice about planned home birth.

All midwives in Canada should provide home birth services to women.

All women under the care of a midwife require unimpeded access to obstetrical and neonatal services and personnel as needed to provide emergent care regardless of intended place of birth.

REFERENCES

- Ackermann-Liebrich U, Voegeli T, Gunter-Witt K, Kunz I, Zullig M, Schindler C, Maurer M. Home versus hospital de-liveries: follow up study of matched pairs for procedures and outcome. Zurich Study Team. BMJ 1996; 313(7068):1313-18.
- Birthplace in England Collaborative Group. Perinatal and maternal outcomes by planned place of birth for healthy women with low risk pregnancies: the Birthplace in England national prospective cohort study. BMJ 2011; 343:d7400.
- 3. De Jonge A, Van der Goes B, Ravelli A, Amelink-Verburga M, Mol B, Nijhuis J, Bennebroek Gravenhorst J, Buitendijk. Perinatal mortality and morbidity in a nation-wide cohort of 529,688 low-risk planned home and hospital births. BJOG 2009; DOI: 10.1111/j.1471-0528.2009.02175.x.
- 4. De Vries R, Buitendijk S. Science, safety and place of birth: lessons from the Netherlands. European Obstet and Gynaecol Suppl 2012; (1):13-17.
- 5. Hafner-Eaton C, Pearce LK. Birth choices, the law, and medicine: Balancing individual freedoms and pro-tection of the public's health. J Health Polit Policy Law 1994: 19:813-35.
- Hutton E, Reitsma A, Kaufman K. Outcomes associated with planned home and planned hospital births in lowrisk women attended by midwives in Ontario, Canada, 2003-2006: A retrospective cohort study. Birth 2009; 36(3):180-89.
- 7. Janssen P, Henderson A, Vedam S. The experience of planned home birth: Views of the first 500 Women. Birth 2009; 36:4:297-304.
- Janssen PA, Lee SK, Ryan EM, Etches DJ, Farquharson DF, Peacock D, Klein MC. Outcomes of planned home births versus planned hospital births after regulation of midwifery in British Columbia. CMAJ 2002; 166(3):315-23.
- 9. Janssen PA, Saxell L, Page LA, Klein MC, Liston RM, Lee SK. Outcomes of planned home births with registered mid-wife versus planned hospital birth with midwife or physician. CMAJ 2009; 181(6):377-83.
- 10. Johnson K, Daviss BA. Outcomes of planned home birth with certified professional midwives: large prospective study in North America. BMJ 2005; 330; 1416.
- 11. Kennare R, Keirse MJ, Tucker GR, Chan AC. Planned home and hospital births in South Australia 1991-2006: differences in outcomes. Med J Aust 2009; 192(2):76-80.
- 12. Lindgren HE; Radestad IJ; Christensson K, Wally-Bystrom K, Hildingsson IM. Perceptions of risk and risk management among 735 women who opted for a home birth. Midwifery 2010; 26(2):163-72.
- 13. Olsen O, Clausen J. Planned hospital birth versus planned home birth. Cochrane Database of Systematic Reviews. September 12, 2012.
- 14. Schroeder E, Petrou E, Patel,N, Hollowell J, Puddi-combe D, Redshaw M, Brocklehurst P. Cost effectiveness of alternative planned places of birth in woman at low risk of complications: Evidence from the Birthplace in England national prospective cohort study. BMJ 2012; 344:e2292.
- 15. Torres Jennifer, De Vries Raymond G. Birthing ethics: What mothers, families, childbirth educators, nurses, and physicians should know about the ethics of childbirth. Peri-nat Educ 2009; 18(1):12–24.
- 16. Vedam S, Schummers L, Stoll K, Fulton C. Home Birth: An annotated guide to the literature. October 2012.
- 17. Vedam, S. Home versus hospital birth: questioning the quality of the evidence on safety. Birth 2003; 30(1):57-
- Wiegers TA, Keirse MJ, van der Zee J, Berghs GA. Out-come of planned home and planned hospital births in low risk pregnancies: prospective study in midwifery practices in the Netherlands. BMJ 1996; 313(7068):1309-13.